


# Methods for the specification and verification of business processes

MPB (6 cfu, 295AA)

Roberto Bruni

<http://www.di.unipi.it/~bruni>

12 - Some Facts



# Object

$$N \vdash \psi$$

We survey  
two connectedness theorems and  
five exchange lemmas

Free Choice Nets (book, optional reading)

<https://www7.in.tum.de/~esparza/bookfc.html>

Two theorems on strong  
connectedness  
(whose proofs are  
optional reading)

# Strong connectedness theorem

**Theorem:** If a weakly connected system is live and bounded then it is strongly connected

(the proof requires some Exchange Lemmas  
that we illustrate later)

# Consequences

If a (weakly-connected) net is not strongly connected

then


It is not live and bounded

If it is live, it is not bounded

If it is bounded, it is not live

# Example

It is now immediate to see that this system  
(weakly connected, not strongly connected)  
cannot be live and bounded  
(it is live but not bounded)



# Exercise

On the basis of the previous observation:

Draw a net that is bounded but not live

Draw a(nother) net that is live but not bounded

Draw a net that is neither live nor bounded

(all nets must be weakly connected)

# Strong connectedness via invariants

**Theorem:** If a weakly connected net has  
a positive S-invariant  $I$  and a positive T-invariant  $J$   
then it is strongly connected



# Consequences

If a (weakly-connected) net is not strongly connected


then

we cannot find (two) positive S- and T-invariants

Five Exchange Lemmas  
(whose proofs are  
optional reading)

# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .  
If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$



# Exchange lemma:

## finite sequences (2)

**Lemma:** Let  $V \subset T$  and  $u \in T \setminus V$ , with  $\bullet u \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma u} M'$  with  $\sigma \in V^*$ , then  $M \xrightarrow{u \sigma} M'$

$$M \xrightarrow{v_1} \xrightarrow{v_2} \cdots \xrightarrow{v_{n-1}} \xrightarrow{v_n} \xrightarrow{u} M'$$

# Exchange lemma:

## finite sequences (2)

**Lemma:** Let  $V \subset T$  and  $u \in T \setminus V$ , with  $\bullet u \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma u} M'$  with  $\sigma \in V^*$ , then  $M \xrightarrow{u \sigma} M'$

$$M \xrightarrow{v_1} \xrightarrow{v_2} \dots \xrightarrow{v_{n-1}} \xrightarrow{u} \xrightarrow{v_n} M'$$

# Exchange lemma:

## finite sequences (2)

**Lemma:** Let  $V \subset T$  and  $u \in T \setminus V$ , with  $\bullet u \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma u} M'$  with  $\sigma \in V^*$ , then  $M \xrightarrow{u \sigma} M'$

$$M \xrightarrow{u} \xrightarrow{v_1} \xrightarrow{v_2} \cdots \xrightarrow{v_{n-1}} \xrightarrow{v_n} M'$$

# More on sequences: projection

**Restriction:** (also extraction / projection)  
given  $T' \subseteq T$  we inductively define  $\sigma|_{T'}$  as:

$$\epsilon|_{T'} = \epsilon \quad (t\sigma)|_{T'} = \begin{cases} t(\sigma|_{T'}) & \text{if } t \in T' \\ \sigma|_{T'} & \text{if } t \notin T' \end{cases}$$

# Example

$$(t_1 t_4 t_7 t_1 t_4 t_7) |_{\{t_1, t_4\}} = t_1 (t_4 t_7 t_1 t_4 t_7) |_{\{t_1, t_4\}}$$

$$= t_1 t_4 (t_7 t_1 t_4 t_7) |_{\{t_1, t_4\}}$$

$$= t_1 t_4 (t_1 t_4 t_7) |_{\{t_1, t_4\}}$$

$$= t_1 t_4 t_1 (t_4 t_7) |_{\{t_1, t_4\}}$$


$$= t_1 t_4 t_1 t_4 (t_7) |_{\{t_1, t_4\}}$$

$$= t_1 t_4 t_1 t_4 (t_7 \epsilon) |_{\{t_1, t_4\}}$$

$$= t_1 t_4 t_1 t_4 (\epsilon) |_{\{t_1, t_4\}}$$

$$= t_1 t_4 t_1 t_4 \epsilon$$

$$= t_1 t_4 t_1 t_4$$






# Exchange lemma:

## finite sequences (3)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma} M'$  with  $\sigma \in (U \cup V)^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V} M'$




# Exchange lemma:

## finite sequences (3)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma} M'$  with  $\sigma \in (U \cup V)^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V} M'$




# Exchange lemma:

## finite sequences (3)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma} M'$  with  $\sigma \in (U \cup V)^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V} M'$




# Exchange lemma:

## finite sequences (3)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .


If  $M \xrightarrow{\sigma} M'$  with  $\sigma \in (U \cup V)^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V} M'$



# Exchange lemma: finite sequences (3)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma} M'$  with  $\sigma \in (U \cup V)^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V} M'$




# Notation $A^\omega$

Given a set  $A$  we denote by  $A^\omega$   
the set of infinite sequences of elements in  $A$ , i.e.:

$$A^\omega = \{ a_1 a_2 \cdots \mid a_1, a_2, \dots \in A \}$$


# Exchange lemma: infinite sequences (4)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V}$



# Exchange lemma: infinite sequences (4)


**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V}$






# Exchange lemma: infinite sequences (4)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V}$



# Exchange lemma: infinite sequences (4)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V}$




# Exchange lemma: infinite sequences (4)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V}$

$$M \quad \underbrace{u_1 u_2 \dots u_{m-1} u_m}_{\sigma|_U} \quad \underbrace{v_1 v_2 \dots v_{n-1} v_n \dots}_{\sigma|_V}$$

# Exchange lemma: infinite sequences (5)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^\omega$ , then  $M \xrightarrow{\sigma|_U}$




# Exchange lemma:

## infinite sequences (5)


**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^\omega$ , then  $M \xrightarrow{\sigma|_U}$




# Exchange lemma: infinite sequences (5)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^\omega$ , then  $M \xrightarrow{\sigma|_U}$



# Exchange lemma: infinite sequences (5)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^\omega$ , then  $M \xrightarrow{\sigma|_U}$



Proofs of exchange  
lemmas  
(optional reading)



# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .  
If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$

Let  $M \xrightarrow{v} K \xrightarrow{u} M'$ .

Clearly  $M' = \underbrace{K - \bullet u + u \bullet}_{K'}$ , with  $K' = K - \bullet u$ .

Since  $\bullet u \cap v \bullet = \emptyset$ , then:  $M'' \xrightarrow{v} K'$  with  $M'' = M - \bullet u$

Therefore:


$$M = M'' + \bullet u \xrightarrow{u} M'' + u \bullet \xrightarrow{v} K' + u \bullet = M'$$

# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$

Let  $M \xrightarrow{v} K \xrightarrow{u} M'$




# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$

Let  $M \xrightarrow{v} K \xrightarrow{u} M'$  and  $K' = K - \bullet u$ .


pre-set of  $u$



# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$



# Exchange lemma:


## finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$

preserved by  
the firing of  $v$

Let  $M \xrightarrow{v} K \xrightarrow{u} M'$  and  $K' = K - \bullet u$ .  
Clearly  $M' = K' + u \bullet$ .




Since  $\bullet u \cap v \bullet = \emptyset$

# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .


If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$



# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$





# Exchange lemma:

## finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$







# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .


If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$



# Exchange lemma: finite sequences (1)

**Lemma:** Let  $u, v \in T$  with  $\bullet u \cap v \bullet = \emptyset$ .

If  $M \xrightarrow{vu} M'$ , then  $M \xrightarrow{uv} M'$



# Exchange lemma: finite sequences (2)

**Lemma:** Let  $V \subset T$  and  $u \in T \setminus V$ , with  $\bullet u \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma u} M'$  with  $\sigma \in V^*$ , then  $M \xrightarrow{u\sigma} M'$

The proof is by induction on the length of  $\sigma$

**base** ( $\sigma = \epsilon$ ): trivially  $M \xrightarrow{u} M'$

**induction** ( $\sigma = \sigma'v$  for some  $\sigma' \in V^*$  and  $v \in V$ ):

Let  $M \xrightarrow{\sigma'} M'' \xrightarrow{vu} M'$ . Note that  $\bullet u \cap v \bullet = \emptyset$

By exchange lemma 1:  $M \xrightarrow{\sigma'} M'' \xrightarrow{uv} M'$ .

Let  $M \xrightarrow{\sigma' u} M''' \xrightarrow{v} M'$ .

By inductive hypothesis:  $M \xrightarrow{u\sigma'} M''' \xrightarrow{v} M'$

Thus,  $M \xrightarrow{u\sigma} M'$

# Exchange lemma: finite sequences (3)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma} M'$  with  $\sigma \in (U \cup V)^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V} M'$

The proof is by induction on the length of  $\sigma|_U$

**base** ( $\sigma|_U = \epsilon$ ): trivially  $\sigma|_V = \sigma$

**induction** ( $\sigma|_U = u\sigma'$  for some  $u \in U$  and  $\sigma' \in U^*$ ):

Let  $M \xrightarrow{\sigma_0} \xrightarrow{u} \xrightarrow{\sigma_1} M'$ , with  $\sigma = \sigma_0 u \sigma_1$  and  $\sigma_0 \in V^*$ .

Note that  $\sigma' = (\sigma_1)|_U$  and  $\bullet u \cap V \bullet = \emptyset$

By exchange lemma 2:  $M \xrightarrow{u} \xrightarrow{\sigma_0} \xrightarrow{\sigma_1} M'$ .

Note that  $(\sigma_0 \sigma_1)|_U = (\sigma_1)|_U = \sigma'$  and  $(\sigma_0 \sigma_1)|_V = \sigma|_V$ .

By inductive hypothesis:  $M \xrightarrow{u} \xrightarrow{\sigma'} \xrightarrow{\sigma|_V} M'$

Since  $\sigma|_U = u\sigma'$ , we conclude that  $M \xrightarrow{\sigma|_U} \xrightarrow{\sigma|_V} M'$

# Exchange lemma: infinite sequences (4)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .  
If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^*$ , then  $M \xrightarrow{\sigma|_U \sigma|_V}$

Let  $\sigma = \sigma' \sigma''$  with  $\sigma'|_U = \sigma|_U$  and  $\sigma''|_V = \sigma''$

(i.e., only transitions in  $V$  appears in  $\sigma''$ ).

Such sequences exist because  $\sigma|_U$  is assumed to be finite.

Let  $M'$  be such that  $M \xrightarrow{\sigma'} M' \xrightarrow{\sigma''}$ .

By Exchange Lemma (3) applied to  $\sigma'$  we have:

$$M \xrightarrow{\sigma'|_U \sigma'|_V} M' \xrightarrow{\sigma''}$$

We conclude by observing that:

$$\sigma|_U = \sigma'|_U \text{ and } \sigma|_V = \sigma'|_V \sigma''$$

# Exchange lemma: infinite sequences (5)

**Lemma:** Let  $U, V \subset T$  and  $U \cap V = \emptyset$ , with  $\bullet U \cap V \bullet = \emptyset$ .

If  $M \xrightarrow{\sigma}$  with  $\sigma \in (U \cup V)^\omega$  and  $\sigma|_U \in U^\omega$ , then  $M \xrightarrow{\sigma|_U}$

To prove that  $M \xrightarrow{\sigma|_U}$  it suffices to show that every finite prefix of  $\sigma|_U$  is enabled at  $M$ .

Take any finite prefix  $\tau'$  of  $\sigma|_U$  and a corresponding finite prefix  $\tau$  of  $\sigma$  such that  $\tau|_U = \tau'$ .

Clearly  $M \xrightarrow{\tau} M'$  for some suitable  $M'$ .

By Exchange Lemma (3), then  $M \xrightarrow{\tau|_U \tau|_V} M'$ , i.e.:  
 $M$  enables  $\tau|_U = \tau'$ .

# Proofs of theorems on strong connectedness (optional reading)

# Strong connectedness theorem

**Theorem:** If a weakly connected system is live and bounded then it is strongly connected

Since the system is live and bounded, by a previous corollary: (see Lecture 11) exists  $M \in [M_0\rangle$  and  $\sigma$  such that  $M \xrightarrow{\sigma} M$  and all transitions in  $T$  occur in  $\sigma$ .

Take any arc  $x \rightarrow y$  in  $F$ :


we need to show that there is a path from  $y$  to  $x$  using arcs of  $F$ .

We distinguish two cases:

1.  $x \in P$  and  $y \in T$
2.  $x \in T$  and  $y \in P$



# Strong connectedness theorem (case 1)



Let  $V = \{v \in T \mid y \rightarrow^* v\}$  and  $U = T \setminus V$ . ( $V$  is the set of transitions reachable from  $y$ )

Note that  $U$  and  $V$  are disjoint and that  $\bullet U \cap V \bullet = \emptyset$ .

(to see this, suppose  $q \in \bullet U \cap V \bullet$  then  $v \rightarrow q \rightarrow u$  for some  $v \in V$  and  $u \in U$ , but then  $u \in V$ , which is impossible because  $U = T \setminus V$ )

By the Exchange Lemma (3), there exists  $M'$  with  $M \xrightarrow{\sigma|_U} M' \xrightarrow{\sigma|_V} M$

We claim that  $M \xrightarrow{\sigma|_V} M$ .

(we want to find a path from  $y$  to  $x$ )

- if  $\sigma|_U = \epsilon$  (i.e.,  $\sigma$  does not contain any transition in  $U$ ), then  $\sigma|_V = \sigma$ .
- otherwise ( $\sigma|_U \neq \epsilon$ ), we can apply the Exchange Lemma (5) to  $M \xrightarrow{\sigma\sigma\cdots}$  to get  $M \xrightarrow{(\sigma\sigma\cdots)|_U} M$ , i.e.,  $M \xrightarrow{\sigma|_U\sigma|_U\cdots} M$ .

Since  $\sigma|_U$  can occur infinitely often from  $M$ , then  $M' \supseteq M$ .


By the Boundedness Lemma  $M' = M$  and  $M \xrightarrow{\sigma|_V} M$ .

Since  $y \in V$ ,  $y$  occurs in  $\sigma|_V$  and  $y \in x \bullet$ , then  $(y$  subtracts a token from  $x)$   
 there must be some transition  $v$  that occurs in  $\sigma|_V$  such that  $v \in \bullet x$ .  $(v$  adds a token to  $x)$

Since  $v \in V$ , there is a path  $y \rightarrow^* v$ .

We can extend this path by the arc  $(v, x)$  to get a path  $y \rightarrow^* x$ .

# Strong connectedness theorem (case 2)



( $U$  is the set of transitions from which  $x$  is reachable)

Let  $U = \{ u \in T \mid u \rightarrow^* x \}$  and  $V = T \setminus U$ .

Note that  $U$  and  $V$  are disjoint and that  $\bullet U \cap V \bullet = \emptyset$ .

(to see this, suppose  $q \in \bullet U \cap V \bullet$  then  $v \rightarrow q \rightarrow u$  for some  $v \in V$  and  $u \in U$ , but then  $v \in U$ , which is impossible because  $V = T \setminus U$ )

(we want to find a path from  $y$  to  $x$ )

By the Exchange Lemma (3), there exists  $M'$  with  $M \xrightarrow{\sigma|_U} M' \xrightarrow{\sigma|_V} M$

By the Exchange Lemma (5) applied to  $M \xrightarrow{\sigma\sigma\cdots}$

we get  $M \xrightarrow{(\sigma\sigma\cdots)|_U} M$ , i.e.,  $M \xrightarrow{\sigma|_U\sigma|_U\cdots} M$ .

Since  $\sigma|_U$  can occur infinitely often from  $M$ , then  $M' \supseteq M$ .

By the Boundedness Lemma  $M' = M$  and  $M \xrightarrow{\sigma|_U} M$ .

Since  $x \in U$ ,  $x$  occurs in  $\sigma|_U$  and  $x \in \bullet y$ , then  $(x$  adds a token to  $y)$   
there must be some transition  $u$  that occurs in  $\sigma|_U$  such that  $u \in y \bullet$ .

$(u$  subtracts a token from  $y)$

Since  $u \in U$ , there is a path  $u \rightarrow^* x$ .

We can extend this path by the arc  $(y, u)$  to get a path  $y \rightarrow^* x$ .


# Strong connectedness via invariants

**Theorem:** If a weakly connected net has  
a positive S-invariant  $I$  and a positive T-invariant  $J$   
then it is strongly connected

Take any arc  $x \rightarrow y$  in  $F$ :  
we need to show that there is a path from  $y$  to  $x$  using arcs of  $F$ .  
We distinguish two cases:

1.  $x \in P$  and  $y \in T$
2.  $x \in T$  and  $y \in P$

# Strong connectedness via invariants: case (1)



Let  $V = \{ v \in T \mid y \rightarrow^* v \}$  and define:

$$J'(t) = \begin{cases} \mathbf{J}(t) & \text{if } t \in V \\ 0 & \text{otherwise} \end{cases} \quad (V \text{ is the set of transitions reachable from } y)$$

(we want to find a path from  $y$  to  $x$ )

Take  $p \in P$ :

- if  $J'(u) = 0$  for all  $u \in \bullet p$ , then:


$$0 = \sum_{u \in \bullet p} J'(u) \leq \sum_{t \in p^\bullet} J'(t)$$

(because  $J'$  has no negative entries).

- otherwise, assume that  $J'(u) = \mathbf{J}(u) > 0$  for some  $u \in \bullet p$ , i.e.,  $y \rightarrow^* u \rightarrow p$ . Then, for any  $t \in p^\bullet$ :  $y \rightarrow^* t$  and  $J'(t) = \mathbf{J}(t) > 0$ . So:

$$0 < \sum_{u \in \bullet p} J'(u) \leq \sum_{u \in \bullet p} \mathbf{J}(u) = \sum_{t \in p^\bullet} \mathbf{J}(t) = \sum_{t \in p^\bullet} J'(t)$$

# Strong connectedness via invariants: case (1)



In both cases:  $\sum_{u \in \bullet p} J'(u) \leq \sum_{t \in p \bullet} J'(t)$

(we want to find a path from y to x)

Then:  $(\mathbf{N} \cdot J')(p) = \sum_{u \in \bullet p} J'(u) - \sum_{t \in p \bullet} J'(t) \leq 0$  for any  $p \in P$ ,

i.e.,  $\mathbf{N} \cdot J'$  has no positive entries.


Since  $\mathbf{I}$  is an S-invariant:  $\mathbf{I} \cdot (\mathbf{N} \cdot J') = (\mathbf{I} \cdot \mathbf{N}) \cdot J' = 0$

and since  $\mathbf{I}$  is positive,  $\mathbf{N} \cdot J' = \mathbf{0}$ , i.e.,  $J'$  is a T-invariant. Hence:

$$\sum_{t \in \bullet x} J'(t) = \sum_{t \in x \bullet} J'(t) \geq J'(y) = \mathbf{J}(y) > 0$$

So there exists  $v \in \bullet x$  with  $J'(v) > 0$ , which means  $v \in V$ , i.e.,  $y \rightarrow^* v$ .  
Since  $v \in \bullet x$ , then  $y \rightarrow^* x$ .

# Strong connectedness via invariants: case (2)




(we want to find a path from  $y$  to  $x$ )

Take  $N' = (T, P, F)$

(i.e., invert the roles of places and transitions).

$N'$



Then,  $\mathbf{N}' = -\mathbf{N}^T$  (where  $\mathbf{N}^T$  is the transposed of  $\mathbf{N}$ )

$\mathbf{I}$  is a positive T-invariant of  $N'$ .

$\mathbf{J}$  is a positive S-invariant of  $N'$ .

By case (1),  $N'$  contains a path from  $y$  to  $x$ .

So,  $N$  contains a path from  $y$  to  $x$ .